

# VLA Observations of H<sub>2</sub>O Masers in the Class 0 Protostar S106

## FIR: Evidence for a 10 AU-Scale Accelerating Jet-like Flow

Ray S. FURUYA

Department of Astronomical Science, The Graduate University for Advanced Studies,  
Nobeyama Radio Observatory

Yoshimi KITAMURA

Institute of Space and Astronautical Science

Masao SAITO

Harvard-Smithsonian Center for Astrophysics

Ryohei KAWABE

Nobeyama Radio Observatory

and

H. Alwyn WOOTTEN

National Radio Astronomy Observatory

Received \_\_\_\_\_; accepted \_\_\_\_\_

MS 39279 — May 25, 1999

## ABSTRACT

We conducted VLA observations at  $0.''06$  resolution of the 22 GHz water masers toward the Class 0 source S106 FIR ( $d = 600$  pc;  $15''$  west of S106-IRS4) on two epochs separated by  $\sim 3$  months. Two compact clusters of the maser spots were found in the center of the submillimeter core of S106 FIR. The separation of the clusters was  $\sim 80$  mas (48 AU) along P.A. =  $70^\circ$  and the size of each cluster was  $\sim 20$  mas  $\times$  10 mas. The western cluster, which had three maser components, was  $8.0$  km s $^{-1}$  blueshifted and the eastern cluster, which has a single component, was  $7.0$  km s $^{-1}$  redshifted with respect to the ambient cloud velocity. Each component was composed of a few spatially localized maser spots and was aligned on a line connecting the clusters. We found relative proper motions of the components with  $\approx 30$  mas year $^{-1}$  (18 AU year $^{-1}$ ) along the line. In addition, a series of single-dish observations show that the maser components drifted with a radial accelerations of  $\sim 1$  km s $^{-1}$  year $^{-1}$ .

These facts indicate that the masers could be excited by a 10 AU-scale jet-like accelerating flow ejected from an assumed protostar located between the two clusters. The outflow size traced by the masers was 50 AU  $\times$  5 AU after correction for an inclination angle of  $10^\circ$  which was derived from the relative proper motions and radial velocities of the maser components. The three-dimensional outflow velocity ranged from 40 to 70 km s $^{-1}$  assuming symmetric proper motions for the blue and red components. Since no distinct CO molecular outflows have been detected so far, we suggest that S106 FIR is an extremely young protostar observed just after the onset of outflowing activity.

*Subject headings:* ISM: jets and outflows — ISM: individual (S106 FIR) — —

stars: formation — masers stars: Class 0 sources — techniques: interferometric

## 1. INTRODUCTION

The formation mechanism of jets and molecular outflows from young stellar objects (YSOs) and their roles in the star formation process are not well understood observationally or theoretically. One of the best observational approaches is to investigate very young protostars which are in an evolutionary stage just after the onset of the outflow activity. André, Ward-Thompson and Barsony (1993) identified such protostars with sources which have not been detected in the near-infrared and which display a blackbody-like spectral energy distribution (SED) which peaks in the submillimeter. They proposed to classify these objects as ‘Class 0’ sources, being in a younger evolutionary stage than that of ‘Class I’ sources defined by Lada (1987). The cold SEDs of Class 0 sources suggest that the bulk of final stellar mass has not yet been accumulated. Class 0 sources are thought to be in their main accretion phase and they are known to have very powerful “jet-like” CO bipolar molecular outflows which are usually observed with typical sizes of  $\sim 0.05 - 0.1$  pc (e.g., Bachiller et al. 1991; Bachiller et al. 1996; Bachiller 1996; Yu and Chernin 1997). In contrast, the CO outflows from Class I sources are poorly collimated and much less powerful (Bontemps et al. 1996). It is believed that the acceleration and collimation of the outflows involve magnetohydrodynamic processes in the vicinity of the central stars ( $r \lesssim 1$  AU). For example, the outflows are thought to be centrifugally accelerated from magnetized Keplerian disks (e.g., Uchida and Shibata 1985; Kudoh and Shibata 1997; Pudritz et al. 1991) or thought to be accelerated from the interaction regions between the stellar magnetospheres and the surrounding disks (e.g., Shu et al. 1988; Najita et al. 1994). Millimeter-interferometer observations with small arrays have extensively investigated the kinematics of gas around Class 0 sources. Their spatial resolutions, however, were at most arcsecond scale (corresponding to a 100 AU scale in nearby star forming regions). It is essential to attain higher spatial resolution – down to subarcsecond-scales – in order to understand outflows in the vicinity of Class 0 sources.

Interferometric imaging of the  $\text{H}_2\text{O}$  maser line at 22 GHz with the VLA and VLBA provides an excellent tool for studying subarcsecond structure in protostellar jets very close to protostars. It has been suggested that the maser lines in low-mass star forming regions are associated with outflows, because the luminosities of the masers correlate with mass-loss rates derived from CO molecular observations (Felli et al. 1992) and with the luminosities of 6 cm free-free continuum emission from ionized, collimated outflows (Wilking et al. 1994; Meehan et al. 1998). Clear evidence for that was obtained by high spatial resolution studies with the VLA-A array (e.g., Chernin 1995) and with the VLBA (e.g., Claussen et al. 1998). These observations revealed that the masers originate behind the shocked gas very close to the central protostars. Interferometric observations of the maser lines have three advantages. First, we can attain high angular resolutions (  $\sim 1\text{-}100$  mas ) and high relative positional accuracies (typically  $\sim 10\%$  of angular resolutions). Second, the maser lines are free from extinction even in deeply embedded sources. Third, we can attain high velocity resolution ( $\lesssim 0.1 \text{ km s}^{-1}$ ) with radio spectroscopy. However, such studies have just begun toward the Class 0/I sources.

We have found that Class 0 sources provide the best targets for studying gas kinematics very close to protostars through observations of the  $\text{H}_2\text{O}$  maser lines. Masers have been detected for 18 of 30 Class 0 sources (detection rate is  $\approx 60\%$ ) in our ongoing multi-epoch survey toward Class 0/I sources with the Nobeyama 45m-telescope (Furuya et al. 1998). On the other hand, we measured a detection rate of only  $\approx 10\%$  for Class I sources. This estimate is based upon data compiled from our survey and previous ones (Wouterloot and Walmsley 1986; Wilking and Claussen 1987; Cessaroni et al. 1988; Comoretto et al. 1990; Terebey et al. 1992; Felli et al. 1992; Palagi et al. 1993; Xiang and Turner 1995; Claussen et al. 1996), although sensitivities and angular and velocity resolutions differ. We interpreted this result as follows. Since Class 0 sources possess a large amount of circumstellar gas, the gas shocked by outflow can easily reach the physical conditions necessary to excite the

maser emission ( $10^6 \text{cm}^{-3} \lesssim n(\text{H}_2) \lesssim 10^9 \text{cm}^{-3}$  with  $T \gtrsim a \text{ few hundred K}$  ; Hollenbach 1997, or  $n(\text{H}_2) \gtrsim 10^9 \text{cm}^{-3}$  with  $T \gtrsim 300 \text{ K}$  ; Elitzur et al. 1989 ). In addition to providing appropriate conditions near the outflow origin, edge-on disks could also enhance the detectability of maser emission by maximizing the path length of velocity coherent gas ( Elmegreen and Morris 1979). In this case, we could investigate kinematics of protostellar disks using the maser probe as shown by Fiebig et al. (1996) in IRAS 00338+6312.

In this paper, we have studied the 22 GHz  $\text{H}_2\text{O}$  maser emission toward the Class 0 source S106 FIR. A compact, bipolar HII region lies near S106 FIR. A distance to S106 molecular cloud is 600 pc (Eiroa, Elsässer and Lahulla 1979; Staude et al. 1982). The exciting star S106-IRS4 (other names are S106-IRS3, S106-IR and S106 PS) whose spectral type is O8V-B0V (Gehrz et al. 1982), is located  $15''$  east of S106 FIR. S106 FIR is a Class 0 source isolated from this bipolar HII region. A dense core with a diameter of 9000 AU was observed around S106 FIR in 450, 800 and  $1100 \mu\text{m}$  continuum emission using the JCMT (Richer et al. 1993), while no emission at  $20 \mu\text{m}$  was detected (Gehrz et al. 1982). The SED of S106 FIR suggests that this source is a Class 0 source. Analysis of the SED implies that the luminosity of S106 FIR is in the range of  $\approx 24$  to  $10^3 L_\odot$  (Richer et al. 1993). It should be noted that a distinct CO molecular outflow has not been detected in S106 FIR, making it unique among all Class 0 sources searched (Hayashi et al. 1993; Bachiller 1996; Furuya et al. 1999), although there is a possibility that wing emission from the S106FIR outflow may be hidden by broad line emission associated with the HII region. The peak of the submillimeter core (see Fig. 4 of Richer et al. 1993) corresponds to the position of the  $\text{H}_2\text{O}$  masers. The maser emission was first detected by Stutzki et al. (1982) and subsequent observations were made by Kawabe (1987) using the Nobeyama Millimeter Array (NMA) in 1987. He found that the masers have a doubly-peaked spectrum with a velocity coverage of  $15 \text{ km s}^{-1}$  and that all the maser spots are concentrated within an area of radius  $0.''3$  (180 AU). At this position, no free-free radio continuum emission has been detected (Felli

et al. 1984; Bally et al. 1983; Hoare et al. 1994). The presence of the H<sub>2</sub>O masers, dense molecular cloud core and the cold SED indicate that S106 FIR may harbor a protostar. In particular, the absence of distinct CO molecular outflows suggests that this protostar is in an extremely early state of star formation.

## 2. OBSERVATIONS

Deep integration synthesis observations of the H<sub>2</sub>O maser line ( $6_{16} - 5_{23}$ ;  $\nu_{\text{rest}} = 22235.077$  MHz) toward S106 FIR were performed with the NRAO<sup>1</sup> Very Large Array (VLA) in its A and BnA configurations on 1996 October 15 and 1997 January 23, respectively. The FWHM of the primary beam was 2.'0 and the system noise temperatures were around 170 K. We observed in the single IF-mode using 128 channels with a total bandwidth of 3.125 MHz. The velocity range covered was 42.1 km s<sup>-1</sup> with a velocity resolution of 0.33 km s<sup>-1</sup> at 22.235 GHz (Table 1). The phase tracking center in the A configuration observations was located at the position reported in the previous NMA measurements (R.A.<sub>1950</sub> = 20<sup>h</sup> 25<sup>m</sup> 32.<sup>s</sup>45, Dec.<sub>1950</sub> = 37° 12' 50."95; Kawabe 1987). The center in the BnA configuration observations was chosen to be at the strongest maser spot in the A configuration observations. The integration time per visibility was 8 seconds, and the field of view (FOV) was limited to be 37'' by a coherency level of 97 % for the longest baseline.

EDITOR: PLACE TABLE 1 HERE.

---

<sup>1</sup>The National Radio Astronomy Observatory (NRAO) is operated by Associated Universities, Inc., under cooperative agreement with the National Science Foundation

The continuum source 2005 + 403 was used as a phase and amplitude calibrator, and was observed for 2 minutes during each 15-minute observing cycle. For the first epoch, 2005 + 403 was also used as a bandpass calibrator by integrating over the run. For the second epoch, 3C454.3 was observed as a bandpass calibrator using 3-minute integration at the beginning and end of the observations. The flux density scale for the first epoch was established using the continuum source 0404 + 768 which was not resolved even in the A configuration. The flux density scale for the second epoch was set by the continuum source 3C286. According to the VLA List of Calibrator Sources, the flux densities of 0404 + 768 and 3C286 at  $\lambda = 1.3$  cm were 1.0 Jy and 2.5 Jy, respectively, and the accuracy is expected to be better than 30%. We made maps of the central regions of  $128 \text{ mas} \times 128 \text{ mas}$  area in the FOV using task IMAGR in the AIPS package with a cell size of 0.5 mas. The resulting synthesized beam sizes were  $63 \text{ mas} \times 61 \text{ mas}$  at P.A. =  $85^\circ$  and  $250 \text{ mas} \times 80 \text{ mas}$  at P.A. =  $89^\circ$  for the first and second epochs, respectively.

We obtained the absolute positions of the strongest maser spots at the velocity channels of  $V_{\text{LSR}} = 6.4 \text{ km s}^{-1}$  ( R.A.<sub>1950</sub> =  $20^h 25^m 32.^s533 \pm 0.^s00015$ , Dec.<sub>1950</sub> =  $37^\circ 12' 50.950'' \pm 0.002''$ ) and  $-13.7 \text{ km s}^{-1}$  ( R.A.<sub>1950</sub> =  $20^h 25^m 32.^s450 \pm 0.^s0006$ , Dec.<sub>1950</sub> =  $37^\circ 12' 50.788'' \pm 0.0025''$ ) for the first and second epochs, respectively, before a self-calibration procedure. The absolute position accuracies of the strongest spots were estimated to be about  $\sigma_{\text{R.A.,Dec}} \approx 4 \text{ mas}$  for the first epoch and  $\sigma_{\text{R.A.}} \approx 14 \text{ mas}$ ,  $\sigma_{\text{Dec}} \approx 5 \text{ mas}$  for the second epoch, respectively, taking account of the typical baseline error of the array ( $|\Delta \mathbf{b}| \approx 1 \text{ cm}$ ) and the angular separation of the phase calibrator from the source ( $|\Delta \mathbf{s}| = 4.9^\circ$ ). These excellent absolute position accuracies result from the use of the close phase calibrator whose coordinates are known with a very high positional accuracy ( $\lesssim 2 \text{ mas}$ ).



In order to obtain the relative positions of the remaining maser spots to the strongest spots at individual channels and to improve dynamic range of images, we have employed a self-calibration procedure using only the phase of the strongest maser spots. The typical rms noise levels in the channel maps after the self-calibration were improved to 10.5 mJy beam<sup>-1</sup> from 28.5 mJy beam<sup>-1</sup> and 9.4 mJy beam<sup>-1</sup> from 22.5 mJy beam<sup>-1</sup> for the first and second epochs, respectively. The relative positional accuracies were determined mainly by the signal-to-noise ratio (S/N ratio), because the bandpass calibration was done within an accuracy of a few degrees across the channels, and because the self-calibration worked very well and the residual systematic errors were very small. The relative positional error of a point source convolved with a synthesized Gaussian beam is given by  $0.45 \theta / (\text{S/N ratio})$  where  $\theta$  is the FWHM of the beam (Reid et al. 1988a;  $\theta = \sqrt{\theta_{\text{maj}} \times \theta_{\text{min}}}$ ). For the first epoch, we use only the data with S/N ratio higher than 7, resulting in a maximum positional error of 5.6 mas (3.4 AU). For the second epoch data, we take the S/N ratio higher than 16 which gives the same maximum positional error as that of the first epoch. These relative positional errors are comparable to those of the absolute ones except for  $\sigma_{\text{R.A.}} \approx 14$  mas of the second epoch.

EDITOR: PLACE TABLE 2 HERE.

We also monitored the masers irregularly in order to investigate the velocity drifts of the maser peaks. The maser emission was observed in 1987 May using the NMA (Kawabe 1987). Single-dish observations were made with the 45m-telescope of Nobeyama Radio Observatory (NRO)<sup>2</sup> on 1996 May 5 and 28. Also 5-minute snapshot observations with

---

<sup>2</sup>Nobeyama Radio Observatory (NRO) is a branch of National Astronomical Observatory, an interuniversity research institute operated by the Ministry of Education, Science, and Culture of Japan

the NRAO, Very Long Baseline Array (VLBA) were conducted on 1996 June 5 as a VLBI Space Observatory Program pre-launch survey (Migenes et al. 1998). The observational parameters are summarized in Table 2.

### 3. RESULTS

#### 3.1. Time Variation of the Overall H<sub>2</sub>O Maser Spectrum

Figure 1 shows the H<sub>2</sub>O maser spectra obtained by the monitoring observations. The intensity scale is shown in Jy unit except for the VLBA snap shot observations because of the uncertainties in the flux calibration. The spectra commonly show three major peaks on the blueshifted side and a single peak on the redshifted side with respect to the ambient cloud velocity ( $V_{\text{sys}} = -1.1 \text{ km s}^{-1}$ ) measured from H<sup>13</sup>CO<sup>+</sup>  $J = 1 - 0$  observations with the NMA (Furuya et al. 1998) and NH<sub>3</sub> ( $J, K$ ) = (2, 2) observations with the VLA (Mangum and Wootten 1994). Distinct maser emission cannot be seen around the ambient cloud velocity. The total radial velocity range of the masers is always from  $V_{\text{LSR}} \approx -15$  to  $12 \text{ km s}^{-1}$ . The blue peaks are commonly seen in the velocity range from  $V_{\text{LSR}} \approx -15$  to  $-2 \text{ km s}^{-1}$ . The red peak is seen from  $V_{\text{LSR}} \approx +2$  to  $+12 \text{ km s}^{-1}$ , although the peak velocity changed from  $V_{\text{LSR}} = +9$  to  $+6 \text{ km s}^{-1}$  in the third quarter of 1996.

EDITOR: PLACE FIGURE 1 HERE.

We found velocity drifts of the spectral peaks in the spectra of Figure 1. The velocities of the three distinct blue peaks drifted together toward the blue, particularly from May 1996 till January 1997. The blue peaks at  $V_{\text{LSR}} = -7.4, -9.4$  and  $-12.1 \text{ km s}^{-1}$  drifted to  $V_{\text{LSR}} = -7.8, -9.8$  and  $-13.7 \text{ km s}^{-1}$ , respectively, during the two epochs of VLA mapping. In addition, we found similar peak velocity drifts on the red side at

$V_{\text{LSR}} = +9 \text{ km s}^{-1}$  from May 1996 until June 1996 and at  $V_{\text{LSR}} = +6 \text{ km s}^{-1}$  during the VLA observations. The red peak at  $V_{\text{LSR}} = 9.2 \text{ km s}^{-1}$  drifted to  $V_{\text{LSR}} = 9.5 \text{ km s}^{-1}$  in about one month and the peak at  $V_{\text{LSR}} = 6.4 \text{ km s}^{-1}$  drifted to  $V_{\text{LSR}} = 6.7 \text{ km s}^{-1}$  in about three months. Here, we fit Gaussian shapes to the spectra in order to determine the peak velocities. These peak velocity drifts can be clearly seen in Figure 2 which shows the peak velocities as a function of the observing date.

The important point which we found from Figure 2 is that individual peaks seem to have their own constant accelerations along the line of sight. For the blue peaks, the accelerations of the velocity drifts are estimated to be  $-2.0$ ,  $-1.1$  and  $-1.1 \text{ km s}^{-1}\text{year}^{-1}$  for the peaks 1, 2 and 3, respectively; and their mean is  $-1.4 \text{ km s}^{-1}\text{year}^{-1}$ . For the red peaks, the peak at  $V_{\text{LSR}} \approx 9 \text{ km s}^{-1}$  showed a velocity drift with an acceleration of  $+3.9 \text{ km s}^{-1}\text{year}^{-1}$  on May 1996. The peak at  $V_{\text{LSR}} \approx 6 \text{ km s}^{-1}$  showed a velocity drift with an acceleration of  $+1.2 \text{ km s}^{-1}\text{year}^{-1}$  during the VLA observations. These velocity drifts are summarized in Table 3.

EDITOR: PLACE FIGURE 2 HERE.

EDITOR: PLACE TABLE 3 HERE.

### 3.2. Spatial Structures of Maser Spots

Figure 3 shows the relative peak positions of the  $\text{H}_2\text{O}$  maser spots in channels of  $0.66 \text{ km s}^{-1}$  resolution, achieved by combining together two adjacent individual channels, for the two epochs of our VLA observations. We plotted maser spots with S/N ratios higher than 7 and 16 for the first and second epochs, respectively. Map origins plotted by

double-circles show the positions of the phase referencing maser spots. A spectrum of the masers integrated over the  $128 \text{ mas} \times 128 \text{ mas}$  area is also shown with a velocity resolution of  $0.33 \text{ km s}^{-1}$  at the upper right corner of each panel. Even at the highest resolution of  $0.33 \text{ km s}^{-1}$ , significant differences cannot be seen in the position and position-velocity maps when compared with Figures 3 and 4 at a resolution of  $0.66 \text{ km s}^{-1}$ . On the other hand,  $0.99 \text{ km s}^{-1}$  resolution exceeds the typical line width of the spectral peaks and causes a serious degradation on the definition of maser components (see §3.4). Thus, we employ a resolution of  $0.66 \text{ km s}^{-1}$  in the maps to avoid the confusion caused by many points.

EDITOR: PLACE FIGURE 3 HERE.

We found that two clusters of maser spots are located in the center of the submillimeter core around S106 FIR and that no maser spots are found outside the  $128 \text{ mas} \times 128 \text{ mas}$  central area. This result is consistent with the previous NMA results (Kawabe 1987). The apparent separation between the intensity-weighted central positions of the two clusters is  $76.2 \text{ mas}$  ( $45.7 \text{ AU}$ ) at  $\text{P.A.} = 72^\circ$  and  $87.5 \text{ mas}$  ( $52.5 \text{ AU}$ ) at  $\text{P.A.} = 70^\circ$  for the first and second epochs, respectively. Each cluster is elongated along a line connecting the two clusters.

### 3.3. Velocity Structures of Maser Spots

Two clusters of the maser spots were found toward the Class 0 source S106 FIR. The western and eastern clusters of maser spots are  $\sim 5 - 13 \text{ km s}^{-1}$  blueshifted and  $\sim 6 - 10 \text{ km s}^{-1}$  redshifted compared to the ambient cloud velocity, respectively. Hereafter, we call them blue and red clusters. The velocity coverages of the blue and red clusters are approximately symmetric with respect to the ambient cloud velocity.

In order to investigate internal velocity structures within each cluster, we show position-velocity maps along two orthogonal lines in Figure 4. The top and bottom panel in Figure 4 are the position-velocity maps along and perpendicular to a line connecting the two clusters. This line is not only parallel to the elongation direction of the clusters but also parallel to proper motions which we will describe in §3.5. We determined the line by  $\chi^2$ -fitting and its position angle was measured to be  $72^\circ \pm 1^\circ$  for October 1996 and  $70^\circ \pm 2^\circ$  for January 1997. We adopt a mean of P.A. =  $71^\circ$ . In Figure 4, filled- and open-circles indicate the maser spots for the first and second epochs, respectively, and the vertical scales of the position-velocity diagrams are the relative radial velocities with respect to the ambient cloud velocity.

EDITOR: PLACE FIGURE 4 HERE.

As is obvious from the top panel of Figure 4, there exist systematic velocity gradients within each cluster along the line at P.A. =  $71^\circ$ , particularly in the blue cluster. The absolute value of velocity increases with increasing the distance from the middle of the two clusters. Mean velocity gradients over the two epochs were  $-0.74 \text{ km s}^{-1} \text{ AU}^{-1}$  for the blue cluster and  $-0.36 \text{ km s}^{-1} \text{ AU}^{-1}$  for the red, determined by  $\chi^2$ -fitting. The velocity gradient for the blue cluster is twice as steep as that of the red cluster. The bottom panel of Figure 4 shows the position-velocity diagram along the line at P.A. =  $-19^\circ$ . All of the maser spots are distributed in a strip of 14 mas (8.4 AU) width centered at the P.A. =  $71^\circ$  line. No clear velocity gradient can be seen along the line at P.A. =  $-19^\circ$  in each cluster.

### 3.4. Identification of Maser Components

In this subsection, we will identify groups of spatially localized maser spots corresponding to the major peaks in the spectra at each epoch. Then we will identify the

maser components across the two epochs.

In order to identify maser components in each epoch, we use the VLA images with velocity resolution of  $0.66 \text{ km s}^{-1}$  (Figure 3) and the spectra of the masers (Figures 1 and 3). In the spectra of the masers, we identified three distinct velocity peaks ( $V_{\text{LSR}} \approx -7, -9$  and  $-12 \text{ km s}^{-1}$ ) on the blue side and a single velocity peak ( $V_{\text{LSR}} \approx 6.0 \text{ km s}^{-1}$ ) on the red side. Here, we call the peaks at  $V_{\text{LSR}} \approx -7 \text{ km s}^{-1}$  as A1 and B1 on the first and second epochs, respectively. Similarly, we label the peaks at  $V_{\text{LSR}} \approx -9$  and  $-12 \text{ km s}^{-1}$  as shown in Tables 4 and 5. In the upper right corner of each panel of Figure 3, the solid bars above the velocity axis show the radial velocity ranges of the three blue peaks and one red peak which we will identify as maser components. In Figure 5, we show maps of the relative positions of maser spots by displaying the velocity ranges of the spectral peaks in pseudocolors. In the upper right corner of each panel, we show the definition of the velocity ranges of the peaks. We found that the maser spots marked by individual colors seem to be spatially localized. Hence, the individual spectral peaks can be identified with these spatial components. We show the relative positions of the components in Tables 4 and 5.

EDITOR: PLACE FIGURE 5 HERE.

Next, we will show correspondence of the maser components between the two epochs. The blue A1, A2 and A3 components on the first epoch are considered to be identical to the blue B1, B2 and B3 components on the second epoch, respectively, from the following kinematical arguments. First, each pair shows a similar velocity range over the two epochs. Second, the peak velocity drifts over  $\sim 250$  days in Figure 1 strongly suggest the presence of three kinematical components on the blueshifted side (i.e., blue 1, 2 and 3), which move with their own constant accelerations (Figure 2). In addition, the red component has a positive acceleration similar to the absolute values of the acceleration of the blue

components, although the red component was observed only over the two epochs. Third, the blue components show similar relative proper motions with respect to the red one, as will be shown in §3.5.

EDITOR: PLACE TABLE 4 HERE.

EDITOR: PLACE TABLE 5 HERE.

### 3.5. Relative Proper Motions of Maser Components

We found an increase in the separations of about 10 mas over the two epochs between the red and three blue components in Figure 5. This fact means that there exist proper motions along the line connecting the two clusters. The absolute positions of the phase referenced maser spots in the two epochs were used to bring self-calibrated relative position maps onto an absolute position basis. This procedure complicates proper motion estimation. Because typical absolute positional errors are  $\sigma'_{\text{R.A.,Dec}} \approx 7$  mas for the first epoch and  $\sigma'_{\text{R.A.}} \approx 15$  mas,  $\sigma'_{\text{Dec}} \approx 8$  mas for the second epoch, respectively, and are comparable to the proper motions, the proper motions are imprecisely determined.

Nevertheless, we can derive relative proper motions from comparison of the relative position maps whose map origins are at the positions of the red spot at  $V_{\text{LSR}} = 6.4 \text{ km s}^{-1}$  for the first epoch and at the red spot at  $V_{\text{LSR}} = 6.7 \text{ km s}^{-1}$  for the second. This superposition is based on our interpretation that the two red spots are physically identical as described in §3.4. We show a magnified H<sub>2</sub>O maser spot map of the blue cluster for the two epochs in Figure 6. Dashed-lines with labels denote the identified maser components. We now find the relative proper motions of the components over the two epochs.

EDITOR: PLACE FIGURE 6 HERE.

The distances between the red component and the three blue components with labels of 1, 2 and 3 increased from 71.7, 73.2 and 83.7 mas to 84.1, 82.1 and 96.3 mas, over the two epochs, respectively. The corresponding proper motions are  $12.4 \pm 5.4$ ,  $8.9 \pm 4.3$  and  $12.6 \pm 3.9$  mas for the blue components 1, 2 and 3, respectively. The proper motion derived for the component 3 was above the  $3\sigma$  level, although the other two were at the  $2\sigma$  level. The directions of the motions are almost parallel to the line connecting the two clusters. Table 6 summarizes the changes of the separations and the relative proper motions. The intensity weighted mean of the proper motions of the blue components relative to the red one is  $10.9 \pm 4.2$  mas ( $6.5 \pm 2.5$  AU) per 100 days. This can be converted to a relative transverse velocity of  $113 \pm 44$  km s<sup>-1</sup> along the plane of the sky.

EDITOR: PLACE TABLE 6 HERE.

## 4. DISCUSSION

We will discuss three possible models for the origin of the two maser clusters: (1) masers associate with a compact jet-like flow, (2) masers originate near each star of a binary system, and (3) masers are generated in tangential parts of a rotating disk. We conclude that the jet-like flow model will explain our results well.

### 4.1. Compact Jet-like Flow Model



#### 4.1.1. Evidence for Compact Outflow and Its Physical Properties

The elongation of the spatial distributions of the maser spots, the presence of the blue and red cluster of maser spots and the relative proper motions of the maser components along the elongation strongly suggest that the maser emission is excited in an expanding jet-like outflow. We assume that the driving source of the outflow, a protostar, is located at the middle of the line connecting the two clusters. This hypothesis is supported by the fact that the spectral profiles of the masers have blue- and redshifted symmetric peaks with respect to the ambient cloud velocity which is thought to be the same as the systemic velocity of the central protostar.

We will discuss three-dimensional geometry of the jet-like flow based on our results. First, we can derive an inclination angle of the outflow out of the plane of the sky from the proper motions and radial velocities of the maser components. The inclination angle  $i$  comes from

$$\tan i = \frac{|V_{\text{mean}} - V_{\text{sys}}|}{\frac{1}{2}\mu} \quad (1)$$

where  $V_{\text{mean}}$  is a mean peak velocity over the two epochs and  $\mu$  is the relative proper motion for each blue component. We obtained  $i = 5.7^\circ, 10.3^\circ$  and  $10.2^\circ$  for the three blueshifted components, respectively (see Table 6); we adopt an intensity weighted mean of  $i = 9^\circ$ . So, the outflow axis of S106 FIR is nearly in the plane of the sky.

Next, we can estimate an outflow velocity range using both the radial velocities and the relative proper motions. Since symmetric proper motions for the blue and red components are suggested by the symmetric maser spectra with respect to the ambient cloud velocity, the outflow velocity is estimated to be one half of the transverse velocity described in §3.5 and ranges from 45 to 65 km s<sup>−1</sup>, corrected for the inclination angle (see Table 6). This derived velocity range is comparable to that reported by Claussen et al. (1998) who measured the proper motions of the H<sub>2</sub>O masers in the Class I source IRAS 05413–0104.

The escape velocity from the protostar of S106 FIR ranges from 10.3 to 20.6 km s<sup>-1</sup> at a distance of 25 AU from the protostar, taking the stellar mass to be 1.5 - 6  $M_{\odot}$  deduced from the observed bolometric luminosity range ( $24L_{\odot} \lesssim L_{\text{bol}} \lesssim 1000L_{\odot}$ ). Hence, the maser spots could not be gravitationally bound to the star: this is consistent with the outflow model.

Furthermore, we estimate the acceleration of the outflow from the radial velocity drifts of the maser components; we assume that the motions of the maser components are limited to lie along the outflow axis. The resulting acceleration ranges from 4 to 15 km s<sup>-1</sup>year<sup>-1</sup> corrected for the inclination angle. We summarize properties of the outflow in Table 7.

EDITOR: PLACE TABLE 7 HERE.

#### 4.1.2. *Formation Mechanism of the Jet-like Flow*

We will discuss two acceleration mechanisms for the jet-like flow. The masers could be accelerated by the stellar wind from the assumed protostar, or by stellar radiation. We conclude that the former mechanism seems most plausible, because the radiation pressure appears insufficient to accelerate the jet-like flow up to the acceleration range estimated in §4.1.1.

First, we will discuss the mechanism whereby the masers are accelerated by the stellar wind which is commonly thought to accelerate the water masers (e.g., Genzel et al. 1981). Here, we consider each identified maser component as a spherical maser cloudlet with a radius of  $a \approx 5$  mas (3AU), which is a typical size of the identified components described in §3.4, for simplicity. Since the acceleration,  $\alpha_W$ , is caused by a pressure gradient over the cloudlet, we have  $\alpha_W = \frac{1}{\rho} \cdot \frac{\Delta P}{a} \sim \frac{1}{\rho} \cdot \frac{P_W}{a}$ , where  $\rho$  is a mass density of the cloudlet

and  $P_W$  is a pressure from the stellar wind. The mass density of the maser cloudlet,  $\rho$  is  $n_{H_2}m_{H_2}$ , where  $n_{H_2}$  is the number density of the cloudlet and  $m_{H_2}$  is the weight of molecular hydrogen. The wind pressure,  $P_W$  at a distance  $r$  from the protostar is given by  $\frac{\dot{M}_w v_w}{4\pi r^2}$ , where  $\dot{M}_w$  and  $v_w$  are the mass loss rate (typically  $\sim 10^{-7} M_\odot \text{year}^{-1}$ ) and wind velocity (typically  $\sim 200 \text{ km s}^{-1}$ ) of protostellar jets, respectively (e.g., Edwards, Ray and Mundt 1994). Thus, we obtain the acceleration from stellar winds as follows

$$\begin{aligned} \alpha_W &\sim \frac{1}{n_{H_2} m_{H_2}} \cdot \frac{1}{a} \cdot \frac{\dot{M}_w v_w}{4\pi r^2} \\ &\sim 7.5 \text{ km s}^{-1} \text{ yr}^{-1} \left( \frac{\dot{M}_w}{10^{-7} M_\odot \text{yr}^{-1}} \right) \left( \frac{v_w}{200 \text{ km s}^{-1}} \right) \left( \frac{25 \text{ AU}}{r} \right)^2 \left( \frac{3 \text{ AU}}{a} \right) \left( \frac{10^8 \text{ cm}^{-3}}{n_{H_2}} \right) \end{aligned} \quad (2)$$

This value seems to be roughly consistent with the acceleration range of  $4 \sim 15 \text{ km s}^{-1} \text{ year}^{-1}$  estimated in §4.1.1. Here, we assumed a typical molecular hydrogen density of  $10^8 \text{ cm}^{-3}$ , necessary to produce maser emission. If we consider smaller cloudlets, down to maser spot sizes of 1 AU (e.g., Reid and Moran 1988b),  $\alpha_W$  becomes larger. However this does not have a serious influence on our discussion. Hence, it is probable that the jet-like flow traced by the masers was mainly accelerated by the stellar wind pressure.

Next, we will discuss the acceleration,  $\alpha_R$ , due to stellar radiation pressure. The acceleration from radiation pressure is given by  $\frac{\kappa L_{\text{bol}}}{4\pi c r^2}$ . If we take the bolometric luminosity,  $L_{\text{bol}}$  to range from 24 to  $1000 L_\odot$  and the mass opacity coefficient,  $\kappa$  to be  $230 \text{ cm}^2 \text{ g}^{-1}$  (a typical value in interstellar medium), then the acceleration from the radiation pressure ranges from  $0.13$  to  $5.2 \text{ km s}^{-1} \text{ year}^{-1}$  at  $r = 25 \text{ AU}$  from the star. Here, we assume that the maser cloudlets are optically thin. If we take an optically thick maser cloudlet,  $\alpha_R$  becomes smaller compared with the optically thin case. This estimated range is smaller than the inclination-corrected acceleration of  $\sim 4 - 15 \text{ km s}^{-1} \text{ year}^{-1}$ . Hence, it seems unlikely that stellar radiation pressure accelerates the jet-like flow.

## 4.2. Other Models for the Origin of the Masers

In this subsection, we discuss the remaining two models for the origin of the masers. First, we will discuss a model in which the masers originate near each star of a protobinary system. Recent infrared and optical observations of pre-main sequence (PMS) stars revealed that the frequency of binary systems in PMS stars is comparable to that in main sequence stars ( $\gtrsim 50\%$ ; e.g., Mathieu 1994; Ghez et al. 1997; Kohler and Leinert 1998; Brandner and Kohler 1998). This fact suggests that binarity must be common among protostars.

In the binary model, the separation of the two maser clusters can be taken as the projected binary separation ( $2a' = 48$  AU). This separation is near the median separation of binaries ( $2a' = 30$  AU from Duquennoy and Mayer 1991). The orbital plane of the binary system is suggested to be perpendicular to the plane of the sky because the proper motions were parallel to the line connecting the two clusters. This is a suitable configuration for the maser amplification. We estimate a rotational velocity of the binary stars in the orbit as  $V_{\text{bin}} = \sqrt{(V_{\text{LSR}} - V_{\text{sys}})^2 + (\frac{1}{2}\mu)^2} \approx 60 \text{ km s}^{-1}$ . A binary phase  $\phi$  measured from the zero projected separation can be estimated from  $\tan \phi = |V_{\text{LSR}} - V_{\text{sys}}|/(\frac{1}{2}\mu)$  for the individual maser components. We adopt a mean value of  $\phi = +10^\circ$ .

The binary model has a serious drawback that the system is not in a gravitationally bound state. For the binary system to be gravitationally bound, the following large enclosed mass is required.

$$M_{\text{enclosed}}(r \leq R_{\text{maser}}) \sim 560 M_{\odot} \left( \frac{a}{140 \text{ AU}} \right) \left( \frac{V_{\text{bin}}}{60 \text{ km s}^{-1}} \right)^2 \quad (3)$$

Here,  $a$  is the half of the binary separation given by  $a = a'/\sin \phi \approx 140$  AU. This enclosed mass is equivalent to a reduced mass of the system. However, the mass is considerably larger than the reduced mass of the binary system expected from the bolometric luminosity range. Therefore, we reject the protobinary model.

Next, we discuss a model of a protostellar disk: a single star is located between the clusters and the masers are excited in tangential parts of a rotating disk around the star. Elmegreen and Morris (1979) originally studied such a model in an ideal case: if there exists enough column density of velocity coherent gas along the line of sight in an edge-on rotating disk, then  $\text{H}_2\text{O}$  masers can be emitted in tangential parts of the disk. The resulting maser spectrum is expected to have a symmetric double-peaked shape with respect to the systemic velocity. The terminal velocity of the spectral peaks corresponds to the rotational velocities of the maser emitting regions of the disk. Water masers originating in disks were found, for the first time, towards the Class I protostar IRAS 00338+6312 with the VLA (Fiebig et al. 1996). The maser profile defined a symmetric double-peaked shape with respect to the ambient cloud velocity. The velocity structure of the maser spots suggests that the  $\text{H}_2\text{O}$  masers are associated with an infalling, rotating disk-like envelope whose radius, traced by the masers, ranges from 10 to 30 AU.

The disk model is not plausible in case of S106 FIR, although the  $\text{H}_2\text{O}$  maser spectra in the source have a symmetric double-peaked shape. In S106 FIR, the disk plane is suggested to be nearly edge-on as mentioned in the binary model, and we can estimate an enclosed mass of the disk and star in the same way as we did for the binary model. The estimated enclosed mass is large, on the order of  $100M_\odot$ , which cannot be reconciled with the bolometric luminosity range. Therefore, we reject the disk model.

#### 4.3. Future Works to Reveal the Nature of S106 FIR

The presence of the  $\text{H}_2\text{O}$  masers and the cold SED suggests that the star formation has just begun in this source. We stress that no well-developed outflows have been reported in S106 FIR (Hayashi et al 1993 and Bachiller 1996). In addition, all of the Class 0 sources except S106 FIR are known to possess distinct CO molecular outflows. These facts suggest

that S106 FIR is at an evolutionary state just after the onset of outflow activity and the formation of well-developed CO molecular outflows usually observed. Do these masers represent a very early state in the development of an outflow from a protostar ?

We have searched for bipolar molecular outflows using the lines of CO  $J=1-0$  and  $\text{HCO}^+$   $J=1-0$  with the NMA and the NRO 45m-telescope, respectively. However, no large-scale molecular outflow has been found (Furuya et al. 1999). On the other hand,  $\text{H}^{13}\text{CO}^+$   $J=1-0$  observations with the NMA and  $\text{NH}_3$  ( $J,K$ )=(1,1) and (2,2) observations with the VLA showed the presence of a dense molecular cloud core around S106 FIR with a mass of  $\sim 1.0 - 1.5M_{\odot}$  and a size of 9000 AU. This dense core is expected to supply molecular gas to a large-scale outflow through interaction with a stellar wind or the compact jet-like outflow.

Our work is being continued as further studies of the proper motions of the  $\text{H}_2\text{O}$  masers with the VLBA. The detailed structure and proper motions of the maser spots will be discussed in a forthcoming paper together with a search for large-scale CO outflow with the NMA. A search for the exact position of the central star(s) by free-free continuum observations will also be attempted. We propose that the central star would be located at the middle of the two clusters ( R.A.<sub>1950</sub> =  $20^h 25^m 32.^s49$ , Dec.<sub>1950</sub> =  $37^\circ 12' 50.''87$ ).

## 5. SUMMARY

High angular resolution VLA observations of the 22 GHz water maser line were conducted toward the Class 0 source S106 FIR. The main findings of this paper are as follows.

1. We found two clusters of the  $\text{H}_2\text{O}$  maser spots with a separation of 50 AU, located at the center of the submillimeter core around the Class 0 protostar S106 FIR. The sizes

of the clusters are  $50 \text{ AU} \times 5 \text{ AU}$ . The western and eastern clusters are  $8.0 \text{ km s}^{-1}$  blueshifted and  $7.0 \text{ km s}^{-1}$  redshifted, respectively, with respect to the ambient cloud velocity.

2. Three maser components were identified in the blue and a single one in the red clusters, respectively. These components are composed of spatially localized maser spots within a  $10 \text{ mas}$  ( $6 \text{ AU}$ ) area, and correspond to the major peaks in the overall spectrum of the  $\text{H}_2\text{O}$  maser line.
3. We determined both the relative proper motions and peak velocity drifts of the maser components. The proper motions are along the line connecting the two clusters and are  $\sim 30 \text{ mas year}^{-1}$ . All the components showed similar peak velocity drifts with radial accelerations of  $\sim 1 \text{ km s}^{-1}\text{year}^{-1}$ .
4. We conclude from our observations that the masers are associated with a highly collimated compact accelerating jet-like flow which could originate from an assumed protostar located at the middle of the line connecting the two clusters.
5. It is possible that S106 FIR is an extremely young protostar in an evolutionary state just after the onset of outflow activity, because of the absence of distinct CO molecular outflows.

The authors gratefully acknowledge an anonymous referee for her very careful reading of the manuscript. Mark J. Claussen and Kevin B. Marvel provided fruitful suggestions as collaborators of our ongoing VLBA study. R. F. acknowledges the kind valuable help by V. Migenes and K. Shibata, during the course of data reduction process with the AIPS. R. F. also appreciates useful discussion and encouragements with M. Tsuboi, T. Omodaka, T. Nakano, S. Kamenno, S. S. Hayashi, T. Kudoh and H. Shinnaga. In addition, authors would

like to thank the staff of NRAO - Socorro and the NRO. M. S. is supported by Smithsonian Postdoctoral Fellowship program.



## REFERENCES

- André, P., Ward-Thompson, D., & Barsony, M. 1993, *ApJ*, 406, 122
- Bachiller, R., Guilloteau, S., Dutrey, A., Planesas, P., & Martin-Pintado, J. 1996, *A&A*, 299, 857
- Bachiller, R., 1996, *ARA&A*, 34, 111
- Bachiller, R., Martin-Pintado, J. & Planesas, P., 1991, *A&A*, 251, 639
- Bally, J., Snell, R. L. & Predmore, R., 1983, *ApJ*, 272, 154
- Bontemps, S., André, P., Terebey, S., & Cabrit, S., 1996. *A&A*, 311, 858
- Brandner & Kohler 1998 *ApJ*, 499, L79
- Cesaroni, R., Palagi, F., Felli, M., Catarzi, M., Comoretto, G., Franco, Di., S., Giovanardi, C., & Palla, F., 1988, *A&AS*, 76, 445
- Chernin, L. M. 1995. *ApJ*, 440, L97
- Claussen, M. J., Wilking, B. A., Benson, P. J., Wootten. H.,A., Myers, P. C., & Terebey, S. 1996, *ApJS*, 106, 111
- Claussen, M, J., Mervel K., B, Wootten, H, A., & Wilking, B, A., 1998, *ApJ*, 507, L79
- Comoretto, G., Palagi, F., Cesaroni, R., et al. 1990, *A&AS*, 84, 179
- Duquennoy, A., & Mayor, M., 1991, *A&A*, 248, 485
- Edwards, S., Ray, T., & Mundt, R. 1994, in *Protostars & Planets III*, eds. E. H. Levy & J. I. Lunine (Tucson : Univ. Arizona Press), 567
- Eiroa, C., Elsässer & Lahulla, J. F. 1979, *A&A*, 74, 89
- Elitzur, M., Hollenbach, D. J., & McKee, C. F, 1989, *ApJ*, 346, 983
- Elmegreen, B., & Morris, M, 1979, *ApJ*, 229, 593

- Felli, M., Staude, H. J., Reddmann, T., Massi, M., Eiroa, C., Hefele, H., Neckel, T., & Panagia, N. 1984. A&A, 135, 261
- Felli, M., Palagi, F., & Tofani, G. 1992. A&A, 255, 293
- Fiebig, D., Duschl, W. J., Menten, K. M., & Tscharnuter, W. M., 1996 A&A, 310, 199
- Furuya, R, S., 1998, unpublished data
- Furuya, R, S., 1999, in preparation
- Gehrz, R. D., Grasdlen, G. L., Castelaz, M., Gullixson, C., Mozurkewich, D., & Hackwell, J., 1982, ApJ, 254, 550
- Ghez, A., Mccarthy, D. W., Patience, J. L., & Beck, T. L. 1997, ApJ, 481, 378
- Genzel, R., Reid, M. J., Moran, J. M., & Downes, D., 1981, ApJ, 244, 884
- Hayashi, S. S., Hasegawa, T., Tanaka, M., Hayashi, M., Aspin, C., Mclean, I. S., Brand, W. J. L., & Gatley, I., 1990. ApJ, 354, L242
- Hollenbach, D., 1997. in *Herbig-Haro Flows and the Birth of Low Mass Stars* , IAU Symposium No. 182, Reipurth, B., and Bertout, C., Editors. 181
- Hoare, G. M., Drew, E. J., Muxlow, B. T., & Davis, J. R., 1994, ApJ, 421, L51
- Kudoh, T., & Shibata, K., 1997, ApJ, 474, 362
- Kawabe, R., 1987 unpublished data
- Kohler, R., & Leinert, C., 1998, A&A, 331, 977
- Lada, C. J. 1987 in IAU Symposium No. 115, Star Forming Regions, ed. M. Peimbert & J. Jugaku (Dordrecht: Kluwer), 1
- Mangum, J. G., & Wootten, A., 1994, BAAS, 26, 1454
- Mathieu, R. D., 1994. ARA&A, 32, 465

- Meehan, L. S., Wilking, B. A., Claussen, M. J., Mundy, L. G., & Wootten, A., 1998. *AJ*, 115, 1599
- Migenes, V., Horiuchi, S., Inoue, M., Edwards, P., Fomalont, E. B., Slysh, V. I. & Val'tis, I. E. 1998, *ASPC* 114, 241,
- Najita, J. R., & Shu, F. H., 1994 *ApJ*, 429, 808
- Palagi, F., Cesaroni, R., Comoretto, G., Felli, M., & Natale, V., 1993, *A&AS*, 101, 153
- Pudritz, R. E. et al. 1991, in *Physics of Star Formation and Early Stellar Evolution*, ed. C. Lada and N. Kylafis (Dordrecht: Kluwer), 539
- Reid, M. J., Schneps, M. H., Moran, J. M., Gwin, C. R., Genzel, R., Downes, D.. & Rönnäng, B. 1988. *ApJ*, 330, 809,
- Reid, M. J., & Moran, J. M.. 1988. in *Galactic and Extragalactic Radio Astronomy* , Verschuur, G. L., and Kellermann, K. I., Editors. (Berlin: Springer), 255
- Richer, J. S., Padman, R., Ward-Thompson, D., Hills, R. E., & Harris, A. I.. 1993. *MNRAS*, 262, 839
- Shu, F. H., Lizano, S., Ruden, S. P., & Najita, J., 1988 *ApJ*, 328, L19
- Staude, H. J., Lenzen, R., Dyck, H. M., & Schmidt, G. D., 1982. *ApJ*, 255, 95
- Stuzki, J., Ungerechts, H., & Winnewisser, G., 1982. *A&A*, 111, 201
- Terebey, S., Vogel, S. N., & Myers, P. C., 1992. *ApJ*, 390, 181
- Uchida, Y., & Shibata, K., 1985. *PASJ*, 37, 515
- Wilking, B. A., Claussen, M. J., Benson, P. J., Myers, P. C., Terebey, S., & Wootten, A., 1994 *ApJ*, 431, L119
- Wilking, B. A., & Claussen, M. J. 1987 *ApJ*, 320, L133
- Wouterloot, J. G. A., & Walmsley, G. M. 1986 *A&A*, 168, 237

Xiang, D., & Turner, B. E. 1995 ApJS, 99, 121

Yu, T., & Chernin, L. M. 1997 ApJ, 479, L63

Table 1: **VLA Mapping Observations of H<sub>2</sub>O Masers toward S106 FIR**

| Epoch       | Config. | Integ. Time<br>(hours) | Synthesized Beam |               | Vel. Coverage<br>(km s <sup>-1</sup> ) | Vel. Resolution<br>(km s <sup>-1</sup> ) |
|-------------|---------|------------------------|------------------|---------------|----------------------------------------|------------------------------------------|
|             |         |                        | Size<br>(mas)    | P.A.<br>(deg) |                                        |                                          |
| 1996 Oct 15 | A       | 3                      | 63 × 61          | 85            | 42.1                                   | 0.3                                      |
| 1997 Jan 23 | BnA     | 2                      | 250 × 80         | 89            | 42.1                                   | 0.3                                      |

---

This manuscript was prepared with the AAS L<sup>A</sup>T<sub>E</sub>X macros v4.0.

Fig. 1.— Spectra of the H<sub>2</sub>O maser emission toward S106 FIR observed with the NMA (1987 May 8), the Nobeyama 45m-telescope (1996 May 5 and 28), the VLBA (1996 June 5) and the VLA (1996 October 15 and 1997 January 23). The intensity scales are shown in Jy unit except for the VLBA observations. The ambient cloud velocity ( $V_{\text{sys}} = -1.1 \text{ km s}^{-1}$ ), measured from H<sup>13</sup>CO<sup>+</sup>  $J = 1 - 0$  and NH<sub>3</sub> ( $J, K$ ) = (2, 2) observations, is also shown. Arrows drawn on the spectra show the drifts of the major peaks ( see in text).

Fig. 2.— Time variations of the velocities of the major peaks of the  $\text{H}_2\text{O}$  maser spectrum in S106 FIR on the redshifted (*upper panel*) and on the blueshifted (*lower panel*) sides. The peak velocities are determined by applying Gaussian profiles to individual peaks.

Fig. 3.— Peak position maps of the  $\text{H}_2\text{O}$  maser spots shown with a velocity resolution of  $0.66 \text{ km s}^{-1}$  with S/N ratio higher than 7 for (a) 1996 October 15 observations and higher than 16 for (b) 1997 January 23. The map origins shown by double-circles are defined by the positions of the phase referencing maser spots at the  $V_{\text{LSR}} = 6.4 \text{ km s}^{-1}$  component on 1996 October and at the  $V_{\text{LSR}} = -13.7 \text{ km s}^{-1}$  component on 1997 January. The spectrum of the masers integrated over the mapping region is also shown at the upper right corner of each panel with a velocity resolution of  $0.33 \text{ km s}^{-1}$ . Solid bars above the velocity axis in the spectra indicate the ranges of the identified velocity components.

Fig. 4.— Upper and lower panels: superposed position-velocity maps of the maser spots on 1996 October 15 observations (filled circles) and on 1997 January 23 (open circles). The upper and lower panels show the position - velocity maps along the line at P.A.=  $71^\circ$  and P.A.=  $-19^\circ$ , respectively. In the ordinate of the position-velocity maps, the ambient cloud velocity ( $V_{\text{sys}} = -1.1 \text{ km s}^{-1}$ ) is subtracted from the radial velocities of the masers spots. Center panel: superposed maps of the maser spots over the two epochs. The two orthogonal lines indicate the lines of P.A.=  $71^\circ$  and P.A.=  $-19^\circ$ , along which we made the position-velocity maps.

Fig. 5.— Peak position maps of identified maser components shown by pseudocolor. The definitions of the components are indicated in the upper right corner of each panel. The map origins are the position of the  $6.4 \text{ km s}^{-1}$  spot for the first epoch, and the position of the  $6.7 \text{ km s}^{-1}$  spot for the second epoch. The phase referencing maser spots in the maps are shown by white-circles.

Fig. 6.— Superposed map of the maser spots at the two epochs for the blue cluster. The positional accuracies are the same as in Figure 3, and the map origins are the same as in Figure 5. Filled and open circles represent 1996 October and 1997 January observations, respectively. Dashed lines with labels indicate the identified maser components.

Table 2: **Summary of H<sub>2</sub>O Maser Observations toward S106 FIR**

| Date        | Telescope         | Beam Size |                 | Vel. Resolution       | Sensitivity <sup>a</sup> |
|-------------|-------------------|-----------|-----------------|-----------------------|--------------------------|
|             |                   | Primary   | Synthesized     |                       |                          |
|             |                   | ( $''$ )  | (mas)           | (km s <sup>-1</sup> ) | (mJy)                    |
| 1987 May 8  | NMA               | 330       | 5000            | 0.26                  | ~100                     |
| 1996 May 5  | NRO 45m           | 74        | ...             | 0.50                  | 180                      |
| 1996 May 28 | NRO 45m           | 74        | ...             | 0.50                  | 240                      |
| 1996 Jun 5  | VLBA <sup>b</sup> | 120       | $\approx 30^c$  | 0.21                  | 220                      |
| 1996 Oct 15 | VLA-A             | 120       | $63 \times 61$  | 0.33                  | 10                       |
| 1997 Jan 23 | VLA-BnA           | 120       | $250 \times 79$ | 0.33                  | 10                       |

<sup>a</sup>Rms noise level per velocity channel.

<sup>b</sup>Performed as a pre-launch survey of VLBI Space Observatory Program (VSOP).

<sup>c</sup>The size is estimated from the fringe spacing of the shortest baseline, between Los Alamos and Pie Town in the VLBA.

Table 3: **Radial Velocity Drifts of H<sub>2</sub>O Maser Peaks in S106 FIR**

| Maser Component | Radial Velocity       |                       | Radial Acceleration |
|-----------------|-----------------------|-----------------------|---------------------|
|                 | 15 Oct 1996           | 23 Jan 1997           |                     |
|                 | (km s <sup>-1</sup> ) | (km s <sup>-1</sup> ) |                     |
| Red             | 6.4                   | 6.7                   | +1.2                |
| Blue 1          | −7.4                  | −7.8                  | −2.0                |
| Blue 2          | −9.4                  | −9.8                  | −1.1                |
| Blue 3          | −12.1                 | −13.7                 | −1.1                |
| Blue (mean)     | ...                   | ...                   | −1.4 <sup>a</sup>   |

---

<sup>a</sup>Intensity weighted mean

Table 4: **Relative Positions of H<sub>2</sub>O Maser Components in S106 FIR on the First Epoch (1996 Oct. 15)**

| Maser Components                       |      | Position Offset <sup>a</sup> |                        | Total Flux Density |
|----------------------------------------|------|------------------------------|------------------------|--------------------|
| Peak Velocity<br>(km s <sup>-1</sup> ) | Name | $\Delta$ R.A.<br>(mas)       | $\Delta$ Dec.<br>(mas) | (Jy)               |
| 6.4 <sup>b</sup>                       | ...  | 0.0                          | 0.0                    | 1.9                |
| -7.4                                   | A1   | $68.9 \pm 1.6$               | $-19.7 \pm 1.6$        | 0.3                |
| -9.4                                   | A2   | $70.5 \pm 0.4$               | $-19.7 \pm 0.4$        | 1.1                |
| -12.1                                  | A3   | $81.3 \pm 0.4$               | $-19.8 \pm 0.4$        | 1.3                |

<sup>a</sup>Intensity weighted mean position

<sup>b</sup>Map origin



Table 5: **Relative Positions of H<sub>2</sub>O Maser Components in S106 FIR on the Second Epoch (1997 Jan. 23)**

| Maser Components                       |      | Position Offset <sup>a</sup> |                        | Total Flux Density |
|----------------------------------------|------|------------------------------|------------------------|--------------------|
| Peak Velocity<br>(km s <sup>-1</sup> ) | Name | $\Delta$ R.A.<br>(mas)       | $\Delta$ Dec.<br>(mas) | (Jy)               |
| 6.7                                    | ...  | $-93.8 \pm 2.8$              | $22.0 \pm 2.7$         | 1.1                |
| -7.8                                   | B1   | $-12.3 \pm 1.6$              | $1.09 \pm 1.0$         | 1.6                |
| -9.8                                   | B2   | $-1.7 \pm 3.4$               | $-13.8 \pm 2.6$        | 2.2                |
| -13.7 <sup>b</sup>                     | B3   | 0.0                          | 0.0                    | 4.0                |

<sup>a</sup>Intensity weighted mean position

<sup>b</sup>Map origin

Table 6: **Relative Proper Motions of H<sub>2</sub>O Maser Components in S106 FIR**

| Name           | Distance from Red Comp. |             | $\mu^a$  | $V_{\text{mean}}$     | $i^b$                                 | $V_{3D}^c$                         |
|----------------|-------------------------|-------------|----------|-----------------------|---------------------------------------|------------------------------------|
|                | 15 Oct 1996             | 23 Jan 1997 |          |                       |                                       |                                    |
|                | (mas)                   | (mas)       | (mas)    | (km s <sup>-1</sup> ) | (degree)                              | (km s <sup>-1</sup> )              |
| Blue 1 (A1→B1) | 71.7±2.3                | 84.1±4.9    | 12.4±5.4 | −7.6                  | 5.7 <sup>+3.6</sup> <sub>−1.9</sub>   | 64 <sup>+1.2</sup> <sub>−5.0</sub> |
| Blue 2 (A2→B2) | 73.2±0.6                | 82.1±4.3    | 8.9±4.3  | −9.6                  | 10.3 <sup>+11.2</sup> <sub>−0.6</sub> | 46 <sup>+1.0</sup> <sub>−4.1</sub> |
| Blue 3 (A3→B3) | 83.7±0.6                | 96.3±3.9    | 12.6±3.9 | −12.9                 | 10.2 <sup>+11.5</sup> <sub>−0.3</sub> | 65 <sup>+1.1</sup> <sub>−5.5</sub> |

<sup>a</sup>Relative proper motions measured from the red component.

<sup>b</sup>Inclination angle defined by  $\arctan\left(\frac{|V_{\text{mean}} - V_{\text{sys}}|}{\mu/2}\right)$ , where  $V_{\text{sys}} = -1.1 \text{ km s}^{-1}$ .

<sup>c</sup>Three-dimensional velocity obtained assuming symmetric proper motions for the blue and red components.

Table 7: **Physical Properties of Compact Jet-like Flow in S106 FIR**

|                                                                            |       |
|----------------------------------------------------------------------------|-------|
| Size (AU)                                                                  | 50×5  |
| P.A. (degrees)                                                             | 70    |
| Inclination (degrees)                                                      | 9     |
| Outflow Velocity <sup>a</sup> (km s <sup>-1</sup> )                        | 45~65 |
| Outflow Acceleration <sup>b</sup> (km s <sup>-1</sup> year <sup>-1</sup> ) | 4~15  |

---

<sup>a</sup>Traced by the masers.

<sup>b</sup>Acceleration, traced by the masers, corrected for the inclination angle of 9° assuming that the motions of the maser components are along the outflow axis.

This figure "Furuya\_Fig1.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9906495v1>

This figure "Furuya\_Fig2.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9906495v1>

This figure "Furuya\_Fig3.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9906495v1>

This figure "Furuya\_Fig4.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9906495v1>

This figure "Furuya\_Fig5.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9906495v1>



This figure "Furuya\_Fig6.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9906495v1>